

Duration of water deprivation in the warm summer months: Effect on reproductive performance of Barbarine rams

S. KHNISSI¹, M. REKIK², N. LASSOUED^{1*}

¹ Laboratoire des Productions Animales et Fourragères, INRAT, Ariana, Tunisia, Université de Carthage.

² International Center for Agricultural Research in the Dry Areas (ICARDA), Amman, Jordan

*Corresponding author: lassoued.narjess@iresa.agrinet.tn

Abstract – Ten mature Barbarine rams were used to study, during hot season, the limit of the tolerance capacity of breed to water deprivation (one, two and three days) on male reproductive performances. Animals were allocated to two groups of five animals each, control (C) and deprived (D) rams. C rams had free access to water while D rams were deprived for 3 days followed by one day of hydration during 4 weeks of mating period (8 cycles of water deprivation). Semen and sexual behavior parameters were recorded every 1st, 2nd and 3rd day of water deprivation during the trial. This study showed an effect of duration of treatment on sperm volume ($P<0.05$) and concentration ($P<0.01$). On the other hand, D rams showed a significant decrease ($p<0.01$) of sexual behavior parameters on the 3rd day of dehydration (anogenital sniffing, flehmen, penis erection, reaction time and libido score). The present study provides a better understanding of the ability of Barbarine breed to endure more than one day dehydration and the response in term of reproductive performances in rams during the mating season which naturally coincides with the summer hot season.

Keywords: Rams, Water deprivation, Semen, Sexual behavior

1. Introduction

In dry areas, water is not available all year-round. Consequently, the animals have to adapt too little water at certain times of the year. Drinking water is an important requirement for livestock and the lack of a sufficient source of water can be a critically limiting factor in animal physiology and productivity (Alamer 2005). Breeds of ruminants native to arid lands graze far away from watering sites sometimes 50 km or more far apart (Silanikove, 1994; Bayer and Feldmann, 2003). In the other hand water intake by sheep is controlled by the capacity of the digestive system. In fact, Casamassima et al, (2016) showed that ovine breeds differ in their respective abilities to respond to water restriction, and experiments on different breeds have yielded conflicting results. Yankasa sheep survive five days without drinking but undergo severe physiological changes (Aganga et al. 1990; Igbokwe 1993); on the other hand, Awassi sheep can withstand more than a month without significant physiological changes while drinking every two days, while watering every five days results in important alterations (Jaber et al. 2004). Mleil et al (2012) showed that recurrent 3 day of water deprivation during last pregnancy - early lactation affect milk production and physiological parameters in Barbarine sheep. Australian Merino sheep could survive 10 days without water (MacFarlane 1964) and the two-horn desert sheep survive for up to 15 days (Farid et al. 1979; Turner 1979). However, Egyptian Barki sheep do not survive for more than three days without water (Farid et al. 1979). In the rams, there is a lack of reports in the literature on the duration of water deprivation and the effect on reproductive parameters. This study was aimed to investigate reproductive parameters, as sperm production and sexual behavior, in semiarid-reared Barbarine sheep in response to different duration of water deprivation. This could lead to better assessment of the relationship between water availability, environmental temperature and reproductive response during mating season.



2. Materials and Methods

2.1. Animals and diets

The trial lasted was carried out on July under natural photoperiod and temperature conditions. Ten sexually experienced mature Barbarine rams were used. All animals had been reared under similar conditions in the research farm of Bou rebiaà of the National Institute of Agricultural Research (INRAT). The station has a semi-arid climate is located 25 km South of the town of Tunis 36°38' N North latitude, 10°07' East longitude, average annual rainfall is 350 mm and represents the mean of 30 years. Animals were allocated to two groups, balanced for live body weight (LBW, 70.5 kg (S.E. 2.7)). Each group of animals was subjected to one of the following two treatments: Treatment (C) receiving water daily ad libitum and Treatment (D) receiving water once every three days. The trial lasted 4 weeks (8 cycles of 3 depriving days followed by one hydration day) Water deprivation in Group D was imposed gradually first for one day, then for 2 days and thereafter for 3 days. Water was offered in buckets (15 L capacity) daily in the morning to Group C, and of the watering days in Group D. Experimental rams were kept in a single barn, well ventilated, with a partition separating the watered and deprived groups. The animals were fed a mixture of straw (1.5 Kg/ram/day) and barley (500g/ram/day) divided into two daily meals (at 08:00 and 19:00 h).

2.2. Experimental measured

All the reproductive parameters (spermiogram and libido test) were recorded on the 1st, 2nd and 3rd days of water deprivation during the 8 cycles. One ejaculate per day for each animal was collected to assess for volume, individual and mass motility, sperm concentration, and the percentage of live spermatozoa. Procedures outlined by Baril et al. (1993) were used to determine these parameters. The rams were trained to ejaculate into an artificial vagina using an estrous ewe as stimulus female for the ejaculation. Libido test was carried out on 6 x 6 m mating pens; rams being individually exposed to an estrous Barbarine ewes. Estrus was induced in these ewes by insertion of progestagen vaginal sponges, as described by Price et al. (1991). Each test lasted 15 minutes and genital sniffing, flehmen reaction intensity, erection of penis, approach to teaser, reaction time and libido score were recorded for each ram (Kilgour and Whale.1980).

The data obtained are presented as (means \pm S.E.M). The statistical analysis was performed according to standard methods (SAS 2005). The analysis of variance (ANOVA) test was used to evaluate the effect of duration (number of days) of water deprivation on the parameters investigated.

3. Results

3.1. Semen parameters

Duration of water deprivation affect significantly sperm volume ($p < 0.05$, table1). In fact, sperm volume decreased significantly from the 2nd day of treatment as compared to the control group ($p < 0.05$). In the other hand we showed a positive relationship between sperm concentration and water deprivation treatment. In fact, D-group showed higher concentration values ($P < 0.01$) compared to C- group on all the trial and mainly in the 3rd day of water deprivation in every cycle. The highest value of D-group sperm concentration as compared to the C-group was recorded in the last cycle of water deprivation of the trial.

3.2. Sexual behavior

A decrease in sexual activity was reflected principally by a decrease in the libido score in water deprivation rams from the second day of water deprivation and becomes significantly ($p < 0.01$, table 2) after 3 days of water deprivation. Indeed, all of the sexual behavior parameters measured have suffered a drop after 3 days of water deprivation except Approach teaser. The reaction time (Time taken by ram to his first mount) was similar in C-rams and in the 1st and the 2nd day of water deprivation but these parameters increased significantly ($p < 0.01$, table 2) at the 3rd day of water deprivation.

Table 1. Means (\pm S.E.M.) values of semen traits at one, two and three days of water deprivation as compared of daily watered group

Parameters	Duration of treatment		S.E.M, Effect DWD
	Daily watering (C)	Water deprivation (D)	
Semen volume (ml)			
1 st day w-d	0.67	0.60	0.03
2 nd day w-d	0.61	0.52	*
3 rd day w-d	0.66	0.46	
Concentration ($\times 10^9$) / ml			
1 st day w-d	7158	7265	182.4
2 nd day w-d	6991	7454	**
3 rd day w-d	6724	8261	
Mass Motility			
1 st day w-d	3.78	3.88	0.1
2 nd day w-d	3.82	3.86	ns
3 rd day w-d	3.71	3.75	
Individual Motility			
1 st day w-d	3	3.37	0.11
2 nd day w-d	3.42	3.21	ns
3 rd day w-d	3.51	3	

ns, not significant; * $P < 0.05$; ** $P < 0.01$. DWD: duration (day) of water deprivation

Table 2. Means (\pm S.E.M.) values of sexual behavior traits at one, two and three days of water deprivation (D) as compared of daily watered group.

Parameters	Duration of treatment		S.E.M, Effect DWD
	Daily watering (C)	water deprivation (D)	
Genital sniffing			
1 st day w-d	1.6	1	0.15 **
2 nd day w-d	1.9	0.2	
3 rd day w-d	2.9	1.5	
Flehmen reaction intensity			
1 st day w-d	1.6	0.3	0.15 **
2 nd day w-d	1.9	1	
3 rd day w-d	2.9	1.5	
Approech to teaser			
1 st day w-d	1.7	1.6	
2 nd day w-d	1.8	1.8	0.11 **
3 rd day w-d	2	1.7	
Erection of penis			
1 st day w-d	2	1.8	0.1
2 nd day w-d	1.9	1.9	ns
3 rd day w-d	2	1.4	
Reaction time (s)			
1 st day w-d	14.4	22.2	1.95 **
2 nd day w-d	12.5	25.1	
3 rd day w-d	12.4	108.9	
Libido score			
1 st day w-d	6.7	6.3	0.21 **
2 nd day w-d	7.3	6.1	
3 rd day w-d	6.6	2.5	

ns, not significant; * $P < 0.05$; ** $P < 0.01$

4. Discussion

Climate change is affecting more and more water reserves especially in semi arid zones. Unavailability of water in addition to high temperatures affects herd productivity at different levels. Reproductive parameters could be an indicator for the welfare and health of animals. Barbarine breed is known by its adaptation capacity to dry areas conditions. Dehydration tolerance can be assessed by the maintain of sufficient energetic state to ensure reproductive activity. The current study revealed significant effect of the water depriving from the third day on the majority of spermatic and sexual behavior parameters.

A number of studies have demonstrated that the spermatogenesis in rams is sensitive to hot temperature (Mahouachi and Khaldi, 1988; Abdel-Hafez, 2002; Marai et al. 2008; Hamilton et al. 2016; Hamilton et al. 2018). Mahouachi and Khaldi (1988) showed, in the Barbarine breed, that high temperature of August decreases semen quality. This effect is added, in our study to water stress and shows that this breed is able to maintain spermatic qualities after two days of dehydration. In fact, water deprivation under dry condition decreased significantly sperm production that was probably due to a decrease of metabolic rate that indicate an energy conservation response (Choshniak et al., 1995). Ejaculation volume depends primarily upon the secretion of seminal plasma rather than concentration which explains the increase on semen cells concentration. On the other hand, there was a tendency to decrease of motility according to the day of collect. This effect could be more pronounced if the dehydration period was extended to more than 60 days, duration of spermatogenesis. The values of mass and individual motility were in the usual values of the breed (Mahouachi and Khaldi, 1988). In the other hand, obtained values in the 3rd day of water deprivation remain in the standards of a spermatic quality which allows a natural mating during the hot season (Khnissi et al, 2015).

Our study showed a drop in sexual activity in deprived rams at the third day of water deprivation as compared to one and two depriving days. Mahouachi and Khaldi (1988) demonstrated in the same breed, an increase of sexual behavior during summer due the seasonality of sexual activity. The observed decrease of sexual behavior was due to dehydration since an imperceptible state of dehydration results in lower of physical capacity. Some authors have suggested that energy has a greater effect on sexual activity because it implies an important physical effort and therefore important energy expenditure (Brown, 1994). In this respect, it is true that water deprived rams uses more energy for thermoregulation. The gradual decrease in libido score and in the time of the first mount in deprived animals depending of the level of water deprivation showed that Barbarine rams can tolerate one and two days of dehydration without an important effect but after three days of water deprivation sexual behavior was dramatically affected and this result is attributed to the depletion of the energy reserves of animals deprived of water in the third day.

5. Conclusion

This study indicates that one or two days of water deprivation can be tolerated by Barbarine rams without an effect on reproductive performances. However, three days of water deprivation in hot season had an effect on some quantitative spermatic parameters, but the semen quality remains in the standards of the breed. In contrary 3 days of water deprivation decreased dramatically sexual behavior of Barbarine ram, which can have a detrimental effect on functioning of the natural mating and consequently an effect on the fertility of the herd.

Acknowledgments

This work was supported by International Atomic Energy Agency (IAEA project TUN5028) and the Tunisian Ministry of Higher Education and Scientific Research. The authors declare that there is no conflict of interest that would prejudice the impartiality of this scientific work.

6. References

- Abdel-Hafez MAM (2002)** Studies on the reproductive performance in sheep. PhD thesis. Faculty of Agriculture, Zagazig University, Zagazig, Egypt
- Aganga A H, Umunna NN, Oyedipe EO, Okoh PN and Aduku AO (1990)** Response to water deprivation by Yankasa ewes under different physiological states. *Small Ruminant Research* 3: 109–115.
- Alamer, M (2005)** Effect of deprivation and season on some biochemical constituents of blood in Awassi and Najdi sheep breeds in Saudi Arabia. *Journal of Animal and Veterinary Advances* 48, 15–20.

- Amir D and Volcani R (1965)** Seasonal fluctuations on the sexual activities of Awassi, Germen Mutton, Merinos, Corriedale, Border Leicester and Dorsett Horn rams. 1. Seasonal changes in semen plasma volume and its fructose and citric acid concentrations. *J. Agric. Sci. (Camb)*, 64(1) : 115-120.
- Baril G, Chemineau P, Cognie Y, Leboeuf B, Orgeur P and Vallet JC (1993)** Collecte et Conservation de la Semence. In : FAO, editor. Manuel de Formation pour L'insémination Artificielle Chez les Ovins et les Caprins. Roma. 7-32.
- Bayer W and Feldmann A (2003)** Diversity of animals adapted to smallholder system. Conservation and Sustainable Use of Agricultural Biodiversity. Available.
- Casamassima D, Vizzarri F, Nardoia M and Palazzo M (2016)** The effect of water-restriction on various physiological variables in intensively reared Lacaune ewes. *Veterinari Medicina*, 61: (11): 623–634.
- Chahal AS, PJS Ratane and O Parshad (1979)** Some physicochemical studies on semen and their interrelationship during different seasons on corriedale rams. *Indian. J. Animal. Sci.*, 49 (6): 433-437.
- Choshniak I, Ben-Kohav N, Taylor CR, Robertshaw D, Barnes RJ, Dobson A, Belkin V and Shkolnik A (1995)** Metabolic adaptations for desert survival in Bedouin goat. *American Journal Physiology* 268: 1101–1110.
- Farid MF, Shawket SM and Abdel-Rahman MHA (1979)** Observations on the nutrition of camels and sheep under stress. In: Camels. IFS Symposium, Sudan. 125–170.
- Galil KAA, Galil AKA (1982a)** Seasonal variation in some characteristics of ejaculated spermatozoa of Sudan Desert sheep in the tropics. *J. Agric. Sci. Camb.*, 99: 35-43.
- Hamilton TRDS, Mendes CM, Signori de Castro L, Monken de Assis P, Perez Siqueira AF, de Carvalho Delgado J, Demarchi Goissis M, Muiño-Blanco T, Álvaro Cebrián-Pérez J, Nichi M, Visintin JA, and Ortiz D'Ávila Assumpção ME (2016)** Evaluation of lasting effects of heat stress on sperm profile and oxidative status of ram semen and epididymal sperm. *Oxidative Medicine and Cellular Longevity*. Volume 2016 12 pages.
- Hamilton TRDS, PSiqueira AF, de Castro LS, Mendes CM, Delgado JDC, De Assis PM, Mesquita LP, Maiorka PC, Nichi M, Goissis MD, Visintin JA, and Ávila Assumpção MEOD (2018)** Effect of Heat Stress on Sperm DNA: Protamine Assessment in Ram Spermatozoa and Testicle. *Oxidative Medicine and Cellular Longevity*. Volume 2018, Article ID 5413056, 14 pages.
- Igbokwe IO (1993)** Haemoconcentration in Yankase sheep exposed to prolonged water deprivation. *Small Ruminant Research* 12: 99–105.
- Jaber LS, Habre A, Rawda N, Abi Said M, Barbour EK and Hamadeh S (2004)** The effect of water restriction on certain physiological parameters in Awassi sheep. *Small Ruminant Research* 54, 115–120.
- KhniSSI S, Lassoued N, Rekik M and Ben Salem H (2014)** Effect of pre-mating water deprivation on Ovarian activity and on reproductive traits of mature Barbarine ewes during summer season. *International Journal of Environmental Science and Development*, 5, 3: 332-335. ISSN 2010-0264.
- KhniSSI S, Lassoued N, Rekik M and Ben Salem H (2015)** Recurrent 3-day cycles of water deprivation for over a month depress sexual behavior but not semen characteristics of adult rams. *Journal of Animal Physiology and Animal Nutrition*. DOI: 10.1111/jpn.12340: 1-8.
- Kilgour R J and Whale R G (1980)** The relation between mating activity of rams in pens and subsequent flock mating performance. *Aus. J. Exp. Agric. Ani. Husb.* 20:5.
- MacFarlane WV (1964)** Terrestrial animals in dry heat: ungulates. In: Dill DB, Adolph EFA, Wilberg CC (eds): *Handbook of Physiology*. American Physiology Society. 509–539.
- Marai IFM, El-Darawany AA, Fadiel A and Abdel-Hafez MAM (2008)** reproductive performance traits as affected by heat stress and its alleviation in sheep. *Tropical and Subtropical Agroecosystems*, 8: 209-234.
- Mleil S, Lassoued N, Ben Salem H and Rekik M (2012)** Effect of water deprivation falling on pregnancy and lactation of mammary secretions on Barbarine ewes and lamb's growth. *European Association of Animal Production Publication* 131: 115–120.
- Price EO, Estep DQ, Wallach SJR and Dally MR (1991)** Sexual performance of rams as determined by maturation and sexual experience. *J. Anim. Sci.* 69: 1047-1052.
- Silanikove N (1994)** The struggle to maintain hydration and osmoregulation in animals experiencing severe dehydration and rapid rehydration: the story of ruminants. *Exper. Physiol.* 79: 281-300.
- Statistical Analysis Sytems (SAS) Institute (2005)** User's guide, version 6.10. SAS Institute, Cary, NC, USA.
- Turner JC (1979)** Osmotic fragility of desert bighorn sheep red blood cells. *Comparative Biochemistry and Physiology* 64: 167–175.