

Long storage and breeder age influences on egg and morphological chick quality

HEDIA NASRI¹, TAHA NAJJAR¹, MONCEF BOUZOUAIA²

¹Laboratory LMMA and UCAR, Department of Animal Sciences, National Agronomic Institute of Tunisia, 43 Avenue Charles Nicolle, Tunis 1082; University of Carthage-Tunisia

²National School of Veterinary Medicine, Ariana, SidiThabet 2020; University of Manouba-Tunisia

*Corresponding author: nasri.hedia@gmail.com

Abstract – A good quality of one-day-old chicks is a crucial hinge between the hatchery and the broiler farm. Storage duration and breeder age are factors that can affect the chick quality. To evaluate this impact, we studied the effect of three storage durations (5, 9 and 23 days) and two breeder ages (31 and 60 weeks) on egg and chick quality.

Long storage (23 days) decreased the egg and the albumen weight and increased the yolk weight, the yolk to albumen ratio ($\Delta = +4.54\%$), the yolk diameter ($\Delta = +2.78$ mm) and the albumen pH ($\Delta = +0.39$, all $p < 0.05$). Long storage (23 days) decreased also, the chick length, the chick weight, the chick length corrected to egg weight (Δ_{23-9} days = -5.49 mm/g), the yolk free body mass (Δ_{23-5} days = -2.15 g) and the Pasgar score (Δ_{5-9} days = -0.27 point; all, $p < 0.05$).

The increase of breeder age increased the egg, the yolk, the albumen and the shell weights as well the yolk to albumen ratio ($\Delta = +9.62\%$) and the yolk diameter ($\Delta = +5.24$ mm) and decreased the shell to egg ratio ($\Delta = -0.99\%$) and the shell thickness ($\Delta = -0.04$ mm; all, $p < 0.05$). The chick length and weight, the yolk free body mass ($\Delta = +6.09$ g), the residual yolk weight and the residual yolk percentage ($\Delta = +2.26\%$) increased also when the breeder age increases. However, the chick length corrected to egg weight ($\Delta = -6.03$ mm/g) and the Pasgar score decreased ($\Delta = -0.2$; all, $p < 0.05$).

Key words: Egg storage, breeder age, egg quality, chick quality.

1. Introduction

In domestic avian species, eggs are stored at cool temperatures until they can be placed into an incubator. Egg storage is a logistical necessity for the hatching egg industry both at the breeder farm and the hatchery (Fasenko, 2007). Hatching eggs can be successfully stored for up to 7 days with little or no effects on hatchability. However, when stored for more than one week, embryonic abnormalities and mortalities increase and general egg quality (yolk membranes, yolk, perivitelline layers) (Decuypere and Bruggeman, 2007) and day old chick quality which quality is the result of events during embryonic development, are affected (Nasri et al., 2019). From the foregoing, it is clear that storage affects embryonic development in different aspects and long storage results in a greater occurrence of poor quality chicks (Boerjan, 2002; Fasenko et al., 2002; Tona et al., 2003; Nasri et al., 2019). The breeder age increase can also negatively affect the egg quality (Lapao et al., 1999), including the internal egg composition (Ulmer-Franco et al., 2010) as well as the chick physical quality (Tona et al., 2004). Old breeder laid heavier eggs than young breeder (Ulmer-Franco et al., 2010). Chicks derived from large eggs have higher body weight and develop faster due to higher availability of nutrients present in the eggs compared to those from small eggs (Lourens et al., 2006). Young breeder lay eggs with thicker eggshells than those from older breeders, and therefore, less oxygen is supplied to the embryos, which may explain their slower metabolism and development (Hamidu et al., 2007). The effect of storage duration is probably interacting with the breeder age and affecting some cited parameters (Tona et al., 2003; Fasenko, 2007; Reijrink et al., 2008). However, published studies of such interactions are limited. The objectives of the experiments described herein were aimed at determining the influence of storage duration, broiler breeder age and their interaction on egg and chick qualities.



2. Materials and methods

2.1. Experimental design

To evaluate the egg quality, the experiment was set up as a 2x2 factorial arrangements with two egg storage durations (5 and 23 days) and two broiler breeder ages (31 and 60 weeks). To evaluate the chick quality, the experiment was set up as a 3x2 factorial arrangements with three egg storage durations (5, 9 and 23 days) and two broiler breeder ages (31 and 60 weeks).

2.2. Eggs and egg storage

In total, 14500 eggs were collected at the same day from three commercial broiler breeder flocks (Arbor Acres), reared in the same conditions as described by the breeding guide (Aviagen, 2018). Breeder flock ages were 31 and 66 weeks. Collected eggs were stored in the hatchery in the same conditions. The storage temperature was 16.5°C and the relative humidity was 70%. The experiment was performed at the Poulina hatchery at Saouef, Tunisia during the months of October and November 2016.

2.3. Egg quality

After 5 and 23 days of storage, 25 eggs per breeder flock age were sampled for egg characteristics (100 eggs in total). Eggs were weighed and thereafter broken onto a flat surface where yolk diameter was determined with an electronic digital caliper (Matrix, Guangdong, China). The albumen was separated from the yolk using a sharp spoon. The yolk was enrolled on absorbent towel to eliminate the rest of albumen and then weighted. Thereafter, the albumen was homogenized separately and its pH was measured immediately with an electronic pH meter (Ohaus, ST2100-F, Parsippany, USA). Shell weight was measured after washing and drying for 24 hours at room temperature. The shell thickness was then determined for the two flocks at 5 days of egg storage using an electronic digital caliper (Matrix, Guangdong, China). For this parameter we evaluated only breeder age impact as the limited impact of storage duration (Samli et al., 2005; Addo et al., 2018). The weight of the albumen was calculated as the difference between the weight of the egg and the weight of the yolk plus the shell. The yolk to albumen ratio was calculated as $((\text{yolk weight} / \text{albumen weight}) * 100)$ and the shell to egg ratio was calculated as $((\text{shell weight} / \text{egg weight}) * 100)$.

2.4. Egg incubation

The rest of the eggs (14400 eggs) were divided into 96 incubation trays of 150 eggs per tray. Each tray contained eggs of one breeder flock and one storage duration. The eggs, placed on incubation trolleys, were stored in the same room. After 5, 9 and 23 days of storage, 16 trays per breeder flock age were incubated. The trays were divided over four incubation trolleys with four trays per breeder age per trolley; the trays being arranged alternatively on trolleys. The four trolleys per storage duration were then placed into two identical Petersime incubators (BioStreamer™, Zulte, Belgium, capacity of 57600 eggs) with 2 trolleys per incubator. All incubators were set at an incubation temperature of (37.78°C). Relative humidity was maintained at an average of 84%. Carbon dioxide level was maintained below 0.85%. Eggs were turned hourly at an angle of 90°.

At 18 days of incubation, all eggs were transferred into hatching crates and moved to hatchers. The same order of trays applied in setters was applied for hatching crates in the hatchers. For each egg storage duration, two Petersime hatchers (BioStreamer™, Zulte, Belgium, capacity of 19200 eggs) were used. All hatchers were set at an incubation temperature (36.67°C). Relative humidity was maintained at an average of 85.50%. Carbon dioxide level was maintained below 1.00%. At 510 hours post-incubation, the chicks were collected from the hatcher.

A number of 40 chicks were chosen randomly per treatment. The chicks length and the Pasgar score (Boerjan, 2002) were determined. Thereafter, 20 chicks per breeder flock age and storage duration were randomly chosen, and killed by cervical dislocation; the body weight and the residual yolk weight were then determined. The live chick body weight was not recorded because the aim was to define the impact of the breeder age and the egg storage duration on the yolk free body mass (YFBM) and residual yolk weight. The YFBM was calculated as the body weight minus the residual yolk weight. The chick length corrected to the egg weight was calculated as $((\text{egg length} / \text{egg weight}) * 100)$ and the body weight corrected to the egg weight was calculated as $((\text{body weight} / \text{egg weight}) * 100)$.

2.5. Statistical Analysis

Data was processed using the statistical software SAS version 9.1 (2004). For egg quality parameters, hatching weight, hatching length and organ weight data, a General linear model (GLM) was performed;

the experimental unit for egg parameters being the egg. The experimental unit for chick weight, chick length and organweight was the chick. The model used for these variables was:

$$Y = \mu + \text{Storage duration} + \text{Breeder age} + \text{Interaction} + e, [1]$$

Where Y= dependent variable, μ = overall mean, Storage duration = storage duration (5, 9 or 23 days), Breeder age = Breeder flock age (31 or 60 weeks of age), Interaction = interaction between storage duration and breeder flock age, e =residual error.

For the Pasgar score a logistic procedure was used to assess the multinomial results; the experimental unit being the chick. For all parameters, the Least Squares Means (LSM) were compared using Bonferroni adjustments for multiple comparisons. Significance was based on $P \leq 0.05$.

3. Results and discussion

3.1. Egg quality

The results concerning the egg quality traits are presented in Table 1. No significant effect of the interaction between the storage duration and the breeder age, was noted in regards to the different studied parameters related to the egg quality ($p > 0.5$).

Table 1: Effect of egg storage duration and broiler breeder age and their interaction on egg characteristics at the start of incubation (LSMeans \pm SEM).

	Egg weight (g)	Yolk weight (g)	Shell weight (g)	Albumen weight (g)	Yolk diameter (mm)	Albumen pH	Yolk to albumen ratio ¹ (%)	Shell to egg ratio ² (%)
Storage								
5	61.51 ^a	18.65	5.40	37.99 ^a	41.68 ^a	8.57 ^a	49.05 ^a	8.87
23	60.12 ^b	19.18	5.46	35.87 ^b	44.46 ^b	8.97 ^b	53.59 ^b	9.13
SEM	0.24	0.24	0.06	0.60	0.2271	0.01	0.78	0.09
Age								
31	54.57 ^a	15.72	5.15 ^a	34.07 ^a	40.45 ^a	8.74 ^a	46.51 ^a	9.49 ^a
60	67.06 ^b	22.12	5.70 ^b	39.79 ^b	45.70 ^b	8.80 ^b	56.13 ^b	8.50 ^b
SEM	0.21	0.20	0.07	0.53	0.2392	0.01	0.92	0.10
Interaction								
5*31	55.66	15.66	5.17	35.34	39.32	8.56	44.60	9.41
5*60	53.48	15.78	5.13	32.80	41.59	8.93	53.49	8.32
23*31	67.37	21.65	5.62	40.65	44.05	8.59	48.41	9.57
23*60	66.76	22.59	5.78	38.94	47.34	9.01	58.76	8.69
SEM	0.23	0.23	0.08	0.67	0.2842	0.02	1.15	0.13
p_value								
Storage	0.02	0.07	0.46	0.01	<.0001	<.0001	0.00	0.05
Age	<.0001	<.0001	<.0001	<.0001	<.0001	0.00	<.0001	<.0001
Interaction	0.18	0.16	0.23	0.58	0.08	0.32	0.58	0.45

a– b: LSM within a column and factor lacking a common superscript differ ($P \leq 0.05$).

¹ Yolk to albumen ratio = (yolk weight/albumen weight)*100

² Shell to egg ratio = (shell weight/egg weight)*100

Long storage decreased the egg ($\Delta = -1.39$ g) and the albumen ($\Delta = -2.12$ g) weight and increased the yolk weight ($\Delta = +0.53$ g) and the yolk to albumen ratio ($\Delta = +4.54\%$). These results agree with the findings of Scott and Silversides (2000). They have noted a decrease in the egg weight from 56.84 g at one day of storage to 56.34 g at 10 days of storage, which was associated with a decrease in the albumen weight from 37.35 g to 35.86 g and an increase in the yolk weight from 13.57 g to 17.47 g. Ahn et al. (1999) reported also a decrease in the albumen percentage from 61.7% to 59.6% and an increase in the yolk percentage from 27.8% to 30.0%, respectively, for 7 and 49 days of storage duration. These phenomena were explained by the loss of water from the egg by evaporation during storage via the porous eggshell and the movement of water from the albumen to the yolk by osmosis (French and Tullet, 1991). Heath (1977) demonstrated also the move of free amino acids from the yolk across the vitelline membrane to the albumen mostly, after 14 days of storage.

Long storage increased the yolk diameter ($\Delta = +2.78$ mm) which may be a result of the weakness of chalazae and vitelline layer that hold the yolk in position and absorb any shocks and jerks to eggs (Alade et al., 2013). Long storage increased also the albumen pH ($\Delta = +0.39$). This result is consistent with Lapao et al. (1999) who noted an increase of the albumen pH from 9.00 to 9.12 points

with an increase of the storage duration from 4 to 8 days. This increase was related to the loss of CO₂ (Stadelman, 1995). As albumen pH increases, the bicarbonate buffering system equilibrium shifts (Heath, 1977). The increase of breeder age increased the egg weight ($\Delta = -1.32$ g), the yolk weight ($\Delta = +6.40$ g), the albumen weight ($\Delta = +5.72$ g), the shell weight ($\Delta = +0.54$ g), the yolk to albumen ratio ($\Delta = +9.62\%$) and the yolk diameter ($\Delta = +5.24$ mm) and decreased the shell to egg ratio ($\Delta = -0.99\%$). This is consistent with Kontecka et al. (2012) who reported an increase in the egg weight by an average of 2.5 g per 5 weeks of breeder age, corresponding to an increase in the yolk and the albumen weights by 1.3 g and 1.1 g, respectively. The amount of shell increased also with breeder age until week 45. However, when considered as a percentage of the egg, the shell and the albumen decreased with the increase of the hen age (Silversides and Scott, 2001), which reveals a higher increase of the yolk weight as compared to that of the albumen. It has been suggested that the increase in the egg size with breeder age is a consequence of a decrease in egg production, which is partly caused by a reduction in the number of follicles reaching the final phase of their rapid growth (Johnson, 2000). Because fewer follicles receive a proportionately greater quantity of yolk, the yolk size of old flock eggs increases at a higher rate as compared to the increase of the albumen size, resulting in a decreased proportion of albumen content (Johnston and Gous, 2007). The increase of breeder age decreased the shell thickness ($\Delta = -0.04$ mm). Roland (1980) associated this decrease to an increase in egg size without a concurrent increase in calcium carbonate deposition. The shell thickness decrease could be one of the reasons of the albumen pH increase ($\Delta = +0.05$ points) with breeder age. In fact, Meijerhof (1994) explain this pH increase by the lower shell quality leading to a higher shell conductance in eggs from older hens that would lead to a more rapid release of carbon dioxide from the egg.

3.2. Chick quality

The results regarding the chick quality traits are presented in Table 2. The interaction between storage duration and breeder age had no influence on the monitored parameters related to chick quality ($p > 0.05$).

Table 2: Effects of egg storage duration, broiler breeder age and their interaction on the quality of one-day-old chicks (LSM \pm SEM).

	Chick length (mm)	Pasgar score	Chick weight (g)	Residual yolk weight (g)	Residual yolk (%)	YFBM ¹ (g)	Body weight corrected to egg weight, (%) ²	Chick length corrected to egg weight (%) ³
Storage								
5		9.42 ^a	42.66 ^a	4.34	10.04	38.32 ^a	69.38	
9	22.71	9.15 ^b	42.61 ^{ab}	4.747	10.98	37.85 ^{ab}	68.99	37.15 ^a
23	18.84	9.35 ^{ab}	40.61 ^b	4.44	10.68	36.17 ^b	67.69	31.68 ^b
SEM	0.056	0.33	0.6002	0.2282	0.4407	0.5016	0.91	0.09
Age								
31	20.50 ^a	9.41 ^a	38.00 ^a	3.59 ^a	9.43 ^a	34.40 ^a	69.31	37.43 ^a
60	21.05 ^b	9.21 ^b	45.92 ^b	5.43 ^b	11.70 ^b	40.49 ^b	68.07	31.39 ^b
SEM	0.055	0.11	0.4921	0.1855	0.3613	0.4113	0.75	0.09
interaction								
5*31		9.52	38.77	3.57	9.14	35.20	69.66	
5*60		9.33	46.55	5.12	10.94	41.44	69.10	
9*31	22.38	9.33	38.33	3.73	9.7	34.56	69.24	40.05
9*60	23.04	8.98	46.89	5.76	12.20	41.13	68.75	34.25
23*31	18.62	9.38	36.92	3.48	9.4	33.44	69.03	34.81
23*60	19.06	9.32	44.30	5.40	11.95	38.90	66.35	28.54
SEM	0.08	0.14	0.86	0.32	0.62	0.70	1.34	0.13
P_value								
Storage	<.0001	0.01	0.03	0.42	0.31	0.01	0.41	<.0001
Age	<.0001	0.01	<.0001	<.0001	<.0001	<.0001	0.25	<.0001
Interaction	0.15	0.40	0.78	0.74	0.81	0.74	0.65	0.07

a– b: LSM within a column and factor lacking a common superscript differ ($P \leq 0.05$).

¹YFBM = chick body weight – residual yolk weight

²Body weight corrected to egg weight, ratio = Chick body weight /egg weight, g/g*100

³Chick length corrected to egg weight ratio = Chick length/egg weight, mm/g*100

Long storage (23 days) decreased the chick length (Δ_{23-9} days = -3.88 mm; $p < 0.0001$), the chick weight (Δ_{23-5} days = -2.05 g; $p = 0.03$), the chick length corrected to egg weight (Δ_{23-9} days = -5.47 mm/g;

$p < 0.0001$) and the yolk free body mass (Δ_{23-5} days = -2.15 g; $p = 0.03$). This is consistent with Goliomytis et al. (2015) who noted the decrease of the chick body weight ($\Delta = -1.24$ g) and the chick length ($\Delta = -0.22$ cm) with the increase of the storage duration from 4 to 16 days of storage. They reported also the absence of significant influence of storage on chick body weight corrected for egg weight prior to setting as well as the chick quality. However, the chick length reported to the egg weight ($\Delta = +0.042$ mm/g) increased. Goliomytis et al. (2015) attributed these results to the lighter eggs before setting resulting from the water loss during storage by evaporation.

The Pasgar score decreased from 5 to 9 days (Δ_{5-9} days = -0.27 point; $p = 0.01$). However, the Pasgar score recorded with 23 days of storage was in the middle. The results concerning the impact of storage on chick quality are conflicting. Many authors reported that the egg storage beyond 7 days was associated with a decline in chick quality (Byng and Nash, 1962; Tona et al., 2003, 2004; Decuyper and Bruggeman, 2007) in terms of physical parameters on the day of hatch such as appearance, activity, and quality of the navel area (Tona et al., 2003). However, Safaa et al. (2013) reported no significant effects on chick quality, in response to the storage duration increase up to 10 days. The same result was reported by Pokhrel et al. (2018) on eggs stored for 7, 14 and 21 days.

The increase of the breeder age increased the chick length (Δ_{31-60} weeks = +0.55 mm; $p < 0.0001$), the chick weight (Δ_{31-60} weeks = +7.91 g; $p < 0.0001$) and the yolk free body mass (Δ_{31-60} weeks = +6.09 g; $p < 0.0001$). These results are consistent with those of Iqbal et al. (2016) who reported an improvement of chick weight and chick length with advancing age of broiler breeder. This improvement was explained by the higher resistance of chicks from old breeder to dehydration upon hatching as compared to smaller chicks from young breeder flocks. However, Nangsuay et al. (2013) did not find any differences in the yolk free body mass when evaluating breeders of different ages (29 and 53 weeks) having similar egg weights. This result confirmed the suggestion of Sklan et al. (2003) that heavier eggs produce heavier chicks and the chick body weight and the yolk free body mass are directly related to the amount of nutrients available in the egg. This is in line with our results concerning the absence of breeder age impact on the chick body weight corrected to the egg weight.

The increase of the breeder age increased the residual yolk weight (Δ_{31-60} weeks = +1.83 g; $p < 0.0001$) and the residual yolk percentage (Δ_{31-60} weeks = +2.26%; $p < 0.0001$). These results agree with the findings of Araújo et al. (2016) who determined a higher yolk sac weight (3.52 g vs. 5.06 g) and relative yolk sac weight (8.64% vs. 10.55%) in chicks from older hens (59 weeks) as compared to young hens (35 weeks). These results can be explained by the higher yolk energy content available in old flock eggs that can probably be related to the yolk nutrient density and especially to the yolk size. These results suggest that older flock hens deposit more yolk and fat in their eggs, particularly in large eggs, than younger flock hens. Consequently, more nutrients will be used for conversion to yolk free body mass with older breeder than younger ones (Nangsuay et al., 2013).

However, the chick length corrected to egg weight (Δ_{31-60} weeks = -6.03 mm/g; $p < 0.0001$) and the Pasgar score decreased (Δ_{31-60} weeks = -0.2; $p = 0.01$) with the increase of the breeder age. Similarly, Tona et al. (2004) observed better physical quality of chicks from 35-week-old hens as compared to those from 45-week-old breeders.

4. Conclusion

Long storage has negative impact on egg quality and on day old chick quality from both, old and young breeders. The manipulation of the storage conditions (Temperature, Humidity ...) could be one of solutions to moderate these negative impacts.

The increase of breeder age increased the vitelline reserve for the embryo by increasing the yolk weight and percentage, which contribute to obtain heavier chicks. However, the one-day-old chick quality based on the Pasgar score decreased with the increase of the breeder age.

5. References

- Ahn DU, Sell J L, Jo C, Chamruspoller TM, Jeffrey M (1999) Effect of Dietary Conjugated Linoleic Acid on the Quality Characteristics of Chicken Eggs During Refrigerated Storage. *Poult. Sci.* 78:922–928. doi: 10.1093/ps/78.6.922

- Alade NK, Usmanand AN, Muhammed AA (2013)** Egg quality Characteristics of two chickens under varying storage conditions and seasons. *Advances in Agriculture, Sciences and Engineering Research*, 3. ISSN: 2276-6723
- Araujo ICSI, Leandro NSMI, Mesquita MAI, Café MBI, Mello HHCI, Gonzales EI (2016)** Effect of incubator type and broiler breeder age on hatchability and chick quality. *Braz. J. Poultry Sci.* 2:17-26. doi: 10.1590/1806-9061-2015-0146
- Aviagen (2018)** Arbor Acres Parent stock handbook. http://es.aviagen.com/assets/Tech_Center/AA_Breeder_ParentStock/AAPSHandBook2018.pdf
- Addo A, Hamidu JA, Ansah AY, Adomako k (2018)** Impact of Egg Storage Duration and Temperature on Egg Quality, Fertility, Hatchability and Chick Quality in Naked Neck Chickens. *Int. J. Poult. Sci.* 17: 175-183. doi: 10.3923/ijps.2018.175.183
- Byng AJ, Nash D (1962)** The effect of egg storage on hatchability. *Br. Poult. Sci.* 3:81-87. doi: 10.1080/00071666208415462
- French NA, Tullet SG (1991)** Variation in the eggs of poultry species. In: *Avian Incubation*. (ed) Tullet S.G., Butterworth-Heinemann Ltd., Cambridge, Great Britain. *Poult. Sci. Symp.* 22, 59-77
- Fasenko GM, Robinson FE, Christensen VL (2002)** How long-term hatching egg storage affects the egg, the embryo and the chick. In: *Practical Aspects of Commercial Incubation in Poultry*. D. C. Deeming, ed. Ratite Conf. Books, Oxford, UK, pp 33-39
- Goliomytis M, Tsipouzian T, Hager-Theodorides A L (2015)** Effects of egg storage on hatchability, chick quality, performance and immuno-competence parameters of broiler chickens. *Poult. Sci.* 94:2257-2265. doi: 10.3382/ps/pev200
- Heath JL (1977)** Chemical and related osmotic changes in egg albumen during storage. *Poult. Sci.* 56 (3): 822-828. doi: 10.3382/ps.0560822
- Iqbal J, Khan S, Mukhtar N, Ahmed T, Pasha RA (2014)** Effects of egg size (weight) and age on hatching performance and chick quality of broiler breeder. *J. Appl. Anim. Res.* doi: 10.1080/09712119.2014.987294
- Johnson AL (2000)** Reproduction in the Female. In: *Sturkie's Avian Physiology* (ed) Whittow, G. C. Academic Press, San Diego, London, Boston, pp. 461-471
- Johnston SA, Gous RM (2007)** Modelling the changes in the proportions of the egg components during a laying cycle. *Br Poult Sci.* 48:347-53. doi: 10.1080/00071660701381134
- Kontecka H, Nowaczewski S, Sierszul MM, Witkiewicz K (2012)** Analysis of changes in egg quality of broiler breeders during the first reproduction period. *Ann. Anim. Sci.* 12:611-622. doi: 10.2478/v10220-012-0051-1
- Lapao C, Gama LT, Chaveiro Soares M (1999)**. Effects of broiler breeder age and length of egg storage on albumen characteristics and hatchability. *Poult. Sci.* 78:640-645. doi: 10.1093/ps/78.5.640
- Meijerhof R (1994)**. The storage of hatching egg. In: *Proceedings of the First International Incubation Conference*. Positive Action Publications Ltd., Utrecht, the Netherlands, 2-5. ISBN 90-74489-13-3
- Molenaar R, Reijrink IAM, Meijerhof R, Van den Brand H (2008)** Relationship between hatchling length and weight on later productive performance in broilers. *World Poult. Sci J.* 64: 599-604. doi: 10.1017/S0043933908000226
- Nangsuay A, Meijerhof R, Ruangpanit Y, Kemp B, van den Brand H (2013)** Energy utilization and heat production of embryos from eggs originating from young and old broiler breeder flocks. *Poult. Sci.* 92:474-482. doi: 10.3382/ps.2012-02643
- Nasri H, van den Brand H, Najjar T, Bouzouaia M (2019)** Egg storage and breeder age impact on egg quality and embryo development. *J. Anim. Physiol. Anim. Nutr.* doi: 10.1111/jpn.13240
- Pokhrel N, Ben-Tal Cohen E, Genin O, Ruzal M, Sela-Donenfeld D, Cinnamon Y (2018)** Effects of storage conditions on hatchability, embryonic survival and cytoarchitectural properties in broiler from young and old flocks. *Poult. Sci.* 97, 1429-1440. doi: 10.3382/ps/pex393
- Roland DA (1980)** Egg Shell Quality: 1. Effect of Dietary Manipulations of Protein, Amino Acids, Energy, and Calcium in Aged Hens on Egg Weight, Shell Weight, Shell Quality, and Egg Production. *Poult. Sci.* 59 (9): 2038-2046. doi: 10.3382/ps.0592047
- Reijrink IAM, Meijerhof R, Kemp B, Van den Brand H (2008)** The chicken embryo and its micro environment during egg storage and early incubation. *Worlds Poult. Sci. J.* 64, 581-598. doi: 10.1017/S0043933908000214

- Stadelman WJ (1995)** The preservation of quality in shell eggs. In: Egg Science and Technology. (ed) Food Products Press, New York, NY, 4th ed. W. J. Stadelman and O. J. Cotterill, pp 67–79
- Scott TA, Silversides FG (2000)** The Effect of Storage and Strain of Hen on Egg Quality. *Poult. Sci.* 79:1725–1729. doi: 10.1093/ps/79.12.1725
- Silversides FG, Scott TA (2001)** Effect of storage and layer age on quality of eggs from two lines of hens. *Poult. Sci.* 80(8):1240-1245. doi: 10.1093/ps/80.8.1240
- Sklan D, Heifetz S, Halevy O (2003)** Heavier chicks at hatch improves marketing body weight by enhancing skeletal muscle growth. *Poult. Sci.* 82:1778–1786. doi: 10.1093/ps/82.11.1778
- Samli HE, Agma A, Senkoylu N (2005)** Effects of Storage Time and Temperature on Egg Quality in Old Laying Hens. *J. Appl. Poult. Res.* 14:548–553. doi: 10.1093/japr/14.3.548
- Safaa HM, Sobhy HN, Elsemary MSA (2013)** Influence of egg storage time on egg quality, hatchability and chick quality traits of commercial and egyptian local broiler breeders. *Aust. J. Basic & Appl. Sci.*, 7(13): 154-163.
- Tona K, Bamelis F, De Ketelaere B, Bruggeman V, Moraes V M B, Buyse J, Onagbesan O, Decuypere E (2003)** Effects of egg storage time on spread of hatch, chick quality, and chick juvenile growth. *Poult. Sci.* 82:736–741. doi: 10.1093/ps/82.5.840
- Tona K, Onagbesan O, De Ketelaere B, Decuypere E, Bruggeman V (2004)** Effects of age of broiler breeders and egg storage on egg quality, hatchability, chick quality, chick weight, and chick posthatch growth to forty-two days. *J. Appl. Poult. Res.* 13:10–18. doi: 10.1093/japr/13.1.10.